

Your healthcare provider has prescribed MORPHABOND ER

Learn how **MORPHABOND ER**
can help manage your pain

MORPHABOND™ ER 
(morphine sulfate) extended-release tablets 
15 mg • 30 mg • 60 mg • 100 mg

What is MORPHABOND ER?

MORPHABOND™ ER (morphine sulfate) extended-release tablets, CII is a strong prescription pain medicine that contains an opioid (narcotic) that is used to manage pain severe enough to require daily, around-the-clock, long-term treatment with an opioid when other pain treatments such as non-opioid pain medicines or immediate-release opioid medicines do not treat your pain well enough or you cannot tolerate them. MORPHABOND ER is a long-acting (extended-release) opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed you are at risk for opioid addiction, abuse and misuse that can lead to death. MORPHABOND ER is not for use to treat pain that is not around-the-clock.

What is important information I should know about MORPHABOND ER?

Get emergency help right away if you take too much MORPHABOND ER (morphine sulfate) extended-release tablets, for oral use, CII (overdose). When you first start taking MORPHABOND ER, when your dose is changed, or if you take too much (overdose), serious or life-threatening breathing problems that can lead to death may occur.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including **BOXED WARNINGS**, and Medication Guide.

CHRONIC PAIN IS PERSONAL

If you suffer from chronic pain, **you are not alone**. Millions of people experience chronic pain, but it affects each person differently.

Because chronic pain is pain lasting more than 3 months, dealing with it can be difficult and frustrating.

Your healthcare professional gave you a prescription for a **medicine to help with your chronic pain**. Your healthcare professional chose **MORPHABOND ER**—one type of medicine used to help with pain

What is important information I should know about MORPHABOND ER? (continued)

Taking MORPHABOND ER with other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death. Never give anyone your MORPHABOND ER. They could die from taking it. Store MORPHABOND ER away from children and in a safe place to prevent stealing or abuse. Selling or giving away MORPHABOND ER is against the law.

Who should not take MORPHABOND ER?

Do not take MORPHABOND ER if you: have severe asthma, trouble breathing, or other lung problems or if you have a bowel blockage or have narrowing of the stomach or intestines.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including **BOXED WARNINGS**, and Medication Guide.

MORPHABOND ER CAN HELP

MORPHABOND ER is a strong prescription pain medicine that contains an opioid (narcotic). It is used to manage pain severe enough to require daily, around-the-clock, long-term treatment with an opioid, when other pain treatments such as non-opioid pain medicines or immediate-release opioid medicines do not treat your pain well enough or you cannot tolerate them.

“ER” means “extended release.” With extended-release opioids, also known as long-acting opioids, the medicine is released more slowly than other pain medicines and is **expected to last for a longer period of time.**



MORPHABOND ER helps your chronic pain by providing relief for **8 to 12 hours after each dose is taken**

Call your healthcare professional if the dose you are taking does not control your pain.

Medicines like MORPHABOND ER can put you at risk for overdose and death. Even if you take your dose correctly as prescribed, you are at risk for opioid addiction, abuse, and misuse that can lead to death.

For more information, visit **MORPHABOND.com**.

HOW TO TAKE MORPHABOND ER

8 or 12^{HRS}



Take your prescribed dose every 8 or 12 hours,
as directed by your healthcare professional.
Do not take more than your prescribed dose. If you
miss a dose, take your next dose at the usual time



Swallow MORPHABOND ER whole.
Do not cut, break, chew, crush, dissolve,
snort, or inject it because this may
cause you to overdose and die



Do not change your dose.

Take MORPHABOND ER exactly as prescribed by
your healthcare professional. Use the lowest dose
possible for the shortest time needed



**Do not stop taking
MORPHABOND ER without talking to
your healthcare professional**

Important things to know about MORPHABOND ER

- Even if you take your dose correctly as prescribed, you are at risk for opioid addiction, abuse, and misuse that can lead to death
- Call your healthcare professional if the dose you are taking does not control your pain
- After you stop taking MORPHABOND ER, flush any unused tablets down the toilet
- Never give anyone your MORPHABOND ER. They could die from taking it
- Store MORPHABOND ER away from children and in a safe place to prevent stealing or abuse
- Selling or giving away MORPHABOND ER is against the law

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including **BOXED WARNINGS**, and Medication Guide.

WHAT TO TELL YOUR HEALTHCARE PROFESSIONAL

Before taking MORPHABOND ER, tell your healthcare professional if you have a history of:

- Head injury
- Seizures
- Liver, kidney, or thyroid problems
- Problems urinating
- Pancreas or gallbladder problems
- Abuse of street or prescription drugs
- Alcohol addiction
- Mental health problems

Tell your healthcare professional if you are:

- Pregnant or planning to become pregnant.** Prolonged use of MORPHABOND ER during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated
- Breastfeeding.** Breastfeeding is not recommended during treatment with MORPHABOND ER. It may harm your baby
- Taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking MORPHABOND ER with certain other medicines can cause serious side effects and could lead to death

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including **BOXED WARNINGS**, and Medication Guide.

MORPHABOND ER CAN HAVE SIDE EFFECTS

The possible side effects of MORPHABOND ER are:

- Constipation
- Nausea
- Sleepiness
- Vomiting
- Tiredness
- Headache
- Dizziness
- Abdominal pain



Call your healthcare professional if you have any of these symptoms and they are severe

Get emergency medical help if you have:

- Trouble breathing
- Shortness of breath
- Fast heartbeat
- Chest pain
- Swelling of your face, tongue, or throat
- Extreme drowsiness
- Lightheadedness when changing positions
- Feeling faint
- Agitation
- High body temperature
- Trouble walking
- Stiff muscles
- Mental changes, such as confusion

These are not all the possible side effects of MORPHABOND ER. Call your healthcare professional for medical advice about side effects.

You may report an adverse event related to Daiichi Sankyo products by calling 1-877-437-7763. You may also report side effects to FDA at 1-800-FDA-1088 or by visiting www.fda.gov/Safety/MedWatch/.

Find **MORPHABOND ER** at your nearest pharmacy

What shouldn't I do while taking **MORPHABOND ER**?

While taking **MORPHABOND ER**, do not:

- ❑ Drive or operate heavy machinery until you know how **MORPHABOND ER** affects you. **MORPHABOND ER** can make you sleepy, dizzy, or lightheaded
- ❑ Drink alcohol or use prescription or over-the-counter medicines that contain alcohol as these may cause you to overdose and die

Call **1-844-438-6111**

to find a pharmacy near you that
carries **MORPHABOND ER**



Before you go to the pharmacy, visit **MORPHABOND.com**
for information about how to save on your prescription.

Eligibility Criteria and Terms and Conditions apply.

Please see additional Important Safety Information
throughout and accompanying full Prescribing Information,
including **BOXED WARNINGS**, and Medication Guide.



Daiichi Sankyo is committed to responsible
use of prescription pain medications

www.CommitmentsinPainCare.com



MORPHABOND ER 
(morphine sulfate) extended-release tablets 
15 mg • 30 mg • 60 mg • 100 mg

©2018 Daiichi Sankyo, Inc.

MORPHABOND ER and the MORPHABOND ER logo are trademarks of INSPIRION DELIVERY SCIENCES LLC.

PP-US-MB-0653 06/18